



XENON100 – The new Results

Marc Schumann *University of Zürich*

CEA Saclay, 22.10.2012

www.physik.uzh.ch/groups/groupbaudis/xenon/




**95% of the
Universe is dark!**



Dark Matter?


Dark Energy????

Dark Matter: (indirect) Evidence





- Particle Dark Matter Candidates:
- WIMP → „WIMP miracle“
 - Axion
 - SuperWIMPs
 - sterile neutrinos
 - WIMPless dark matter
 - Gravitino



Dark Matter Search



Indirect
Detection




Production
@Collider



Direct WIMP Search

Elastic Scattering of
WIMPs off target nuclei
→ nuclear recoil



Recoil Energy:

$$E_r = \frac{|\vec{q}|^2}{2m_N} = \frac{\mu^2 v^2}{m_N} (1 - \cos \theta) \sim \mathcal{O}(10 \text{ keV})$$

Event Rate:

$$R \propto N \frac{\rho_\chi}{m_\chi} \langle \sigma_{\chi-N} \rangle$$

Detector

Local DM Density

Physics

N number of target nuclei
 ρ_χ/m_χ local WIMP density
 $\langle \sigma \rangle$ velocity-averaged scatt. X-section

Direct WIMP Search




Summary: Tiny Rates

$$R < 0.01 \text{ evt/kg/day}$$

$$E_R < 100 \text{ keV}$$

How to build a WIMP detector?

- large total mass, high A ✓ for Xe
- low energy threshold ✓ for Xe
- ultra low background ✓ for Xe
- good background discrimination ✓ for Xe




Backgrounds

Experimental Sensitivity



without background: $\propto (\text{mt})^{-1}$
 with background: $\propto (\text{mt})^{-1/2}$

Background Sources

environment: U, Th chains, K




- γ and β decays (electronic recoil)
- alphas no big problem for LXe (technology dependent)
- neutrons from (α, n) and sf in rocks and detector parts
- neutrons from cosmic ray muons





Why WIMP Search with Xenon?

- efficient, fast scintillator (178nm)
- high mass number $A \sim 131$:
 - SI: high WIMP rate @ low threshold
- high $Z=54$, high $\rho \sim 3$ kg/l:
 - self shielding, compact detector
- SD: 50% odd isotopes
 - allows further characterization after detection by testing only SI or SD
- no long lived Xe isotopes
 - Kr can be removed to ppt level
- "easy" cryogenics @ -100°C
- scalability to larger detectors
- in dual-phase TPC:
 - good background discrimination





Dual Phase TPC





Dual Phase TPC

3d Vertex Reconstruction




Signal/Background Discrimination




The XENON program

XENON: A phased WIMP search program




XENON
R&D


2005-2007:
XENON10



2008-2013:
XENON100



2010-2017:
XENON1T



Columbia



Rice



UCLA



U Zürich



Coimbra



LNGS



Mainz



SJTU



Bologna



MPIK



NIKHEF



Purdue



Subatech



Münster



WIS



XENON Collaboration



XENON Collaboration Meeting, LNGS, October 2012



XENON100

Astropart. Phys. 35, 573 (2012)




161 kg LXe, 62 kg in target
242 1" x1" PMTs


Laboratori Nazionali del Gran Sasso



XENON100 Background




- 30 kg fiducial mass
- active LXe veto not used for this plot
- exploit anti-correlation between light and charge for better ER-energy scale




Xenon keVee-Scale not precisely known below 9 keVee

Measured Background in good agreement with MC prediction.


At low energies: Lowest background ever achieved in a Dark Matter Experiment!





Low Energy Response to ER




Compton Scatter Measurement at UZH
indicates that LXe „sees“ electronic recoil
interactions around ~ 2.3 keV (also with field!!!)



we are currently analyzing data
taken at the XENON100 field
→ needed for modulation analysis



(spin-independent) WIMP Limit 2011




XENON100 sets the most sensitive limit over a large WIMP mass range
Challenges the CoGeNT, DAMA, CRESST-II signals as being due to light mass WIMPs

PRL 107, 131302 (2011)
already cited 408x

Limit derived with
Profile Likelihood method
PRD 84, 052003 (2011)

OUTDATED!!!




XENON100 – New results of 2012

arXiv:1207.5988, accepted by PRL

Data Taking


Data Collection



Data taking over 13 months
from Feb 28, 2011 to March 31, 2012
→ full annual cycle

3 interruptions for maintenance
224.56 live days of dark matter data



Stability




To our knowledge, no large LXe detector has ever been operated under such stable conditions for that long

Improvements

- Exposure more than doubled
- Lower threshold
 $S2 > 150 \text{ PE}$, $S1 > 3 \text{ PE}$ (6.6 keV ν)
- Lower Background
- Much more calibration data
35x more ER calibration in ROI
AmBe before and after run
- Higher LXe purity \rightarrow smaller corrections




ER/NR Discrimination



Discrimination comparable to previous runs:
 ~99.5% ER rejection @ 50% NR acceptance


Total ER Background



ER Background:
(with active veto)



$(5.3 \pm 0.6) \times 10^{-3}$ events/keV/kg/day
→ before discrimination

Data Analysis: All data





More information on XENON100 data analysis in [arXiv:1207.3458](https://arxiv.org/abs/1207.3458)


Basic Quality Cuts



Single Scatter Selection





Threshold and Fiducial Volume




Consistency Cuts


S2 Width:




Select Energy Range



ER Rejection




WIMPs are Nuclear Recoil-like



Profile Likelihood Method

PRD 84, 052003 (2011)



background + WIMP signal
(100 GeV/c² at 10⁻⁴⁴ cm², 13 events)

observed Signal

expected background

Construct Likelihood function:

need good understanding of background
("background model")
→ but this is required by any
low background experiment
(regardless of the type of analysis)

$$\mathcal{L} = \mathcal{L}_1(\sigma, N_b, \epsilon_s, \epsilon_b, \mathcal{L}_{\text{eff}}, v_{\text{esc}}; m_\chi) \times \mathcal{L}_2(\epsilon_s) \times \mathcal{L}_3(\epsilon_b) \times \mathcal{L}_4(\mathcal{L}_{\text{eff}})$$

NR scale measurement


"NR like" sideband measurement

dark matter measurement


"ER like" sideband measurement

Cuts and Acceptance

S2 Threshold acceptance
(to be applied before S1 smearing)



independent fluctuations



details: [arXiv:1207.3458](https://arxiv.org/abs/1207.3458)

Cuts Acceptance



NR acceptance
(benchmark region only)

Nuclear Recoil Energy Scale

- WIMPs interact with Xe nucleus
 - nuclear recoil (nr) scintillation (β and γ 's produce electronic recoils)
- absolute measurement of nr scintillation yield is difficult
 - measure relative to ^{57}Co (122keV)
- relative scintillation efficiency L_{eff} :

$$\mathcal{L}_{\text{eff}}(E_{\text{nr}}) = \frac{\text{LY}(E_{\text{nr}})}{\text{LY}(E_{\text{ee}} = 122 \text{ keV})}$$

measurement principle:



most recent measurements:

- *Plante et al., PRC 84, 045805 (2011)*
- △ *Manzur et al., PRC 81, 025808 (2010)*

for discussion of possible systematic errors see
A. Manalaysay, arXiv:1007.3746

Background Prediction

Neutron background:

- $(\alpha, n) + sf$ and muon induced neutrons
- MC simulation using the exact XENON100 geometry and measured contaminations

Expect: $(0.17 +0.12 -0.07)$ events



ER background:

- γ activity of the detector and shield
- intrinsic radioactivity in the LXe
(→ considerably lowered this run)
- use ER calibration to model background by scaling it to the observable DM data

Expect: (0.79 ± 0.16) events

Sum: (1.0 ± 0.2) events

The same background model is implemented in the PL analysis




During ...



... and after ...



... Unblinding





(1.0 ± 0.2) events expected
2 events observed

- 26.4% probability that background fluctuated to 2 events
- PL analysis cannot reject the background only hypothesis


No significant excess due to a signal seen in XENON100 data.

Events in Benchmark Region


- visual inspection:
valid waveforms
- at 7.1 keVr and 7.8 keVr
both events between
3 and 4 PE
- rather low wrt the
NR calibration data
- no low S2/S2-events
below threshold




The new XENON100 Limit




No Impact of L_{eff} below 3 keVr




The new XENON100 Limit



What XENON100 sees...




A light mass WIMP...




$$m_\chi = 8 \text{ GeV/c}^2 \quad \sigma = 3.0 \times 10^{-41} \text{ cm}^2$$

A CRESST-like signal...




$$m_\chi = 25 \text{ GeV/c}^2 \quad \sigma = 1.6 \times 10^{-42} \text{ cm}^2$$

What XENON100 excludes...



$$m_\chi = 50 \text{ GeV}/c^2 \quad \sigma = 3.0 \times 10^{-45} \text{ cm}^2$$

What could the Events be?




Reminder:

Background is modeled using ER calibration data from Co60 and Th232

This data shows an increased probability for anomalous leakage below ~8 PE

Background prediction depends on the information which is put into the model


Sensitivity to single electrons




Relaxing the S2 threshold condition ($\text{S2}>150$ PE)
leads to a band of events at very low S2/S1 (below signal range)

- can the 2 events be in the tail of this band???
- further studies are required
- aim: quantify and put into background model for the next run


XENON: What next?



How do we get there?



The next step: XENON1T




- 3t LXe ("1m³ detector")
1t fiducial mass → 20x larger
- 100x lower background
(~10 cm self shielding,
low radioactivity components)
- background goal: <1 evt in 2 years




Low Radioactivity
Photon Detectors
(3", Total ~250)


low radioactivity
stainless steel
cryostat
(or copper)



The next step: XENON1T




- 3t LXe ("1m³ detector")
1t fiducial mass → 20x larger
- 100x lower background
(~10 cm self shielding,
low radioactivity components)
- background goal: <1 evt in 2 years
- Timeline: 2010 – 2017
- start construction early 2013






XENON1T @ LNGS



XENON1T
@ LNGS (Hall B)
→ 4.8 m radius water shield
acting as active muon veto

- Proposal and TDR submitted to LNGS
- Approved by INFN end of April 2011
- Approved by NSF (US) May 2012



The new WIMP Landscape

