



Discussion on 10<sup>th</sup> May

Due on 17<sup>th</sup> May

**Exercise 1** *Kronig-Penney Model*

With this exercise, we will solve the Kronig-Penney model in small steps. This model starts with the assumption that the potential that the electron feels near the atomic positions can be approximated by a  $\delta$ -function. The periodic potential can thus be written as  $U(x) = Aa \sum_s \delta(x - sa)$  (in one dimension) where  $A$  is a constant and  $a$  is the lattice spacing.

(a) Show that in the Fourier series  $U(x) = \sum_G U_G e^{iGx}$  we get for the coefficients  $U_G = A$ .

(b) Let  $\psi(k)$  be the Fourier coefficients of the electron wave function. Use the central equation (derived in the lecture or found in text books and on the web) to show that

$$(\lambda_k - \epsilon)\psi(k) + A \sum_n \psi(k - 2\pi n/a) = 0 \quad (1)$$

where  $\lambda_k = \hbar^2 k^2 / 2m$ .

(c) Now we are interested in solving with respect to  $\epsilon$ . In this context it is convenient to define  $f(k) = \sum_n \psi(k - 2\pi n/a)$ . Show that (Caution:  $n$  has two meanings):

$$\psi(k) = -\frac{(2mA/\hbar^2)f(k)}{k^2 - (2m\epsilon/\hbar^2)} \quad (2)$$

$$f(k) = f(k - 2\pi n/a) \quad (3)$$

$$\psi(k - 2\pi n/a) = -\frac{(2mA/\hbar^2)f(k)}{(k - 2\pi n/a)^2 - (2m\epsilon/\hbar^2)} \quad (4)$$

(d) Now let's sum ( $\sum_n$ ) on both sides of equation 4 found above and show that:

$$\hbar^2/2mA = -\sum_n [(k - 2\pi n/a)^2 - (2m\epsilon/\hbar^2)]^{-1} \quad (5)$$

Notice how with these simple operations we got rid of the Fourier coefficients of the wave function  $\psi$ .

(e) Our goal is still to solve with respect to  $\epsilon$ . Let's define  $K^2 = 2m\epsilon/\hbar^2$  and remember that  $\cot(x) = \sum_n \frac{1}{n\pi+x}$ . Show that:

$$\frac{\hbar^2}{2mA} = \frac{a^2 \sin(Ka)}{2Ka(\cos(ka) - \cos(Ka))} \quad (6)$$

Hint: It is helpful to use a partial fraction decomposition of equation 5. The following trigonometric identities might be useful:

$$\cot x - \cot y = \frac{\sin(y - x)}{\sin x \sin y} \quad (7)$$

$$2 \sin x \sin y = \cos(x - y) - \cos(x + y) \quad (8)$$

(f) What is the energy of the lowest energy band at  $k = 0$  if we assume  $P = \frac{mAa^2}{\hbar^2} \ll 1$ ? (Hint: how big can  $Ka$  be? If it is small, you can expand in  $Ka$ .)

(g) Evaluate the band gap at the zone boundary  $k = \pi/a$  (we still assume  $P \ll 1$ ).

(h) Now let us set  $P = 3\pi/2$ . Plot the dispersion relation from 0 to  $4\pi/a$ . Plot also the same dispersion in the reduced zone scheme (all bands folded into first zone). (Hint: You have to solve equation 6 numerically. If you do not see any energy gaps, you did something wrong. Furthermore, the dispersion relation should look somewhat parabolic. Pay attention to the starting values.)